Post Fire Seeding

Erosion and Weed Control

Washington State (east of Cascade Mts.)

For more information contact your local NRCS office



The following are broadcast seeding PLS rates. If seeding with a drill, reduce by half.

PLS = **Pure Live Seed** = %Purity x %Germination (required tests)

% purity and % germination will be on the seed tags (all seed should come with a seed tag)

Native species (n) - use plant materials with germplasm from the Inland Northwest

	Broadcast Seeding Rate		
	Seeds/lb.	Rate	Seeds
Species*	approx.	lbs/ac	per. sq. ft.
Rangeland (native mix)			
(n) Thickspike wheatgrass (Schwendimar, Bannock)	135,000	6	19
(n) Big bluegrass 'Sherman'	925,000	1	21
(n) Sandberg bluegrass (Mt Home, 'Reliable')	1,000,000	1	23
(n) Snake River wheatgrass ('Secar' or 'Discovery')	139,000	6	19
or (n) Bluebunch wheatgrass			
Total		14	82
Rangeland (non-native mix)			
Pubescent wheatgrass	85,000	8	16
Intermediate wheatgrass	85,000	8	16
White Dutch clover	776,000	1	18
Total		17	49
Forest (native mix)			
(n) Mountain brome (<i>may substitute blue wildrye</i>)	65,000	6	9
(n) Slender wheatgrass (may substitute blue wildrye)	135,000	4	12
(n) bluebunch wheatgrass (may sub Snake River wheatgrass)	135,000	8	25
(n) Junegrass (Koeleria macrantha)	2,300,000	0.5	26
Total		18.5	73
Forest (non-native mix)			
Orchardgrass	540,000	3	37
Timothy	1,200,000	1	28
(n) Mountain brome	65,000	8	12
Total		12	77
Emergency stabilization			
Winter wheat	14,000	15-25	5 - 8
or - Regreen (sterile wheat x wheatgrass hybrid)	12,000	15-25	4 - 7
or - Quickguard (sterile hybrid triticale)	13,000	15-25	4.5 - 7.5
*May use local Source Identified seed	as well		
Broadcast seeding should be done in the fall o		r.	
For pasture mix recomendations contact your loc			